



Game Planning for Infectious Diseases



Presented by: Cherie L. Drenzek, DVM, MS

State Epidemiologist

Date: 11/12/2014

We Protect Lives.





NFL Game Planning

- Scouting Reports
- Stats
- Film Study
- Opponents' Tendencies
- Neutralize Strengths
- Exploit Weaknesses


GOAL: WIN!



NFL Game Planning



NFL Game Planning



Infectious Disease Game Planning


GOAL:

STOP THE SPREAD OF INFECTION


How? PPE, handwashing, vaccines, antibiotic treatment, disinfection, decontamination, food recall, restaurant closure, and others...

Infectious Disease Game Planning: How do we know how to stop transmission?

- Know your Opponent
 - Know the Chain of Infection
 - Understand the Natural History of a Disease
 - Understand the Epidemiology of a Disease
- Exploit Weaknesses



Game Planning First Step: Tailgating






Infectious Disease Game Planning Know Your Opponent: Microbes

- What is a Microbe?
 - Our unseen friends and foes
 - Friends produce wine, foes cause human (or animal) infection and illness
 - They are everywhere!
 - Types:
 - Bacteria (Salmonella, TB, E.coli, MRSA)
 - Viruses (Ebola, norovirus)
 - Yeast and Fungi (Candida, Exserohilum)
 - Protozoa (Cyclospora, Cryptosporidium)




Infectious Disease Game Planning Know the Chain of Infection



Opponents: Infectious Agents


- Found in nature in <u>reservoirs</u> (its natural habitat)
- Reservoirs can be people (pertussis, norovirus), animals (Ebola, Salmonella, E.coli O157:H7), or inanimate matter like soil (botulism, tetanus, anthrax)
- Microbes survive and reproduce in the reservoir. Sometimes the microbe makes the reservoir sick, sometimes not.



Chain of Infection: How do microbes exit the reservoir and infect someone else?

Modes of Transmission


- Many ways! Many definitions! Many levels!
- Direct vs Indirect
- Vehicles (Food, Fomites)
- Vector (Mosquito, Tick)
- Person-to-Person
- Airborne (Measles, TB), Bloodborne (HIV, Hepatitis B)
 Fluid-borne (Ebola), Foodborne (Salmonella),
 Waterborne (Cryptospordium)
- Zoonotic (Ebola, MERS, Rabies, Q fever)





Chain of Infection: How long does it take?

- Incubation period
 - Time from exposure to the microbe to the development of symptoms
- Examples: Flu 1-4 days, Ebola 2-21 days,
 TB 3-12 weeks, rabies 2 months-1 year
- Contagious or Shedding Period may precede symptoms
- Knowing both **key** to risk assessment and control




Patterns (Or, the Epidemiology of our Opponents)



- It's a Small World After All: Emerging infectious diseases are only a plane, boat, ship, or bus ride away
- In 2014, the world shared these infectious diseases: Influenza A H7N9, Chikungunya, MERS, and Ebola
- For foodborne outbreaks: widely distributed contaminated food products; many fresh produce vehicles
- Prevalence of antimicrobial resistance (MRSA, MDR-TB, CRE) and healthcare-associated infections


How Does This Help Us Break the Chain?



- Control at any link in the chain
- Some easier
- Reservoir harder
- ★Transmission/entry = PPE, disinfection, handwashing
- Susceptible host = vaccination, isolation, quarantine
- Also focus on epidemiology: early detection among travelers

Some Examples of ID Game Plans






Bacterial Agents: Salmonella

- Reservoir: Animals (poultry, cattle, amphibians), Humans (typhi), SEROTYPE-specific
- Incubation period: 1-3 days
- Clinical symptoms: diarrhea, fever, abdominal cramps, SEVERE, death
- Epidemiology/patterns:
 - Restaurants, environment, PFGE clusters
- Associated vehicles:
 - Eggs, chicken, fresh produce (melon, tomatoes, sprouts)
- Spread: Cross-Contamination from Food to Environment to Person



Bacterial Agents: Salmonella



Viral Agents: MERS Coronavirus

- Reservoir: Camels?
- Incubation period: 2-14 days
- Clinical symptoms: Severe acute respiratory illness, fever, cough, death.
- Epidemiology/patterns:
 - Travelers to Middle East (21 countries), healthcare spread, some close contact spread
- Spread: Direct contact, droplet?
- Control: WHO says people with diabetes, kidney failure, or chronic lung disease and weakened immune systems should:
 - Avoid contact with camels
 - Do not drink raw camel milk or raw camel urine
 - Do not eat undercooked camel meat








Bacterial Agents: *E.coli* O157:H7

- **Reservoir**: Cattle
- **Incubation period**: 1-8 days
- Clinical symptoms: Severe/bloody diarrhea, abdominal pain, vomiting, HUS, death.
- Epidemiology/patterns:
 - Restaurants, events, fairs, water/recreational water
- Associated vehicles:
 - Undercooked ground beef, fresh produce (apples, sprouts), water
- Spread: Cattle feces to beef or to produce via water or manure

Bacterial Agents: *E. coli* O157:H7 Where's The Beef?



EVD Natural History: Transmission

Ebola is spread through direct contact

How do you get the Ebola virus?


Direct contact with:

- Body fluids of a person who is sick with or has died from Ebola.

 (blood, vomit, pee, poop, sweat, semen, spit, other fluids)
- Objects contaminated with the virus (needles, medical equipment)
- Infected animals (by contact with blood or fluids or infected meat)

Is NOT Airborne

Epi Patterns: Ebola Early Case Recognition



For more information on how to care for a person under investigation for Ebola, please visit: http://www.cdc.gov/ebola

e Protect Lives.

Toxin Producers: Staphylococcus aureus/MRSA

- **Reservoir**: Humans (nose, throat, boils, piercings); Animals
- **Incubation period**: Varies
- Clinical symptoms: Skin infection, boils, purulent, "spider bite"
- Epi patterns:
 - Athletic Teams (Tampa Bay Buccaneers), healthcare settings
- Associated vehicles: towels, razors, shared equipment, whirlpool
- Spread: skin-to-skin direct contact, indirect via vehicles above
- Control: Keep wounds covered, handwashing, no sharing





Parting Words

- We rarely mentioned "You Know Who"--"Eboldemort" –don't worry--more to come!
- Etiologic natural history is the foundation of our game plan to defeat our opponent microbes and the keys to their control.
- Infectious disease preparedness has multi-purpose benefits; partnerships always the key to overall improved community health.
- And if something just doesn't seem right...

How to Contact Epidemiology (24/7)

Office/Epi On Call (8-5): 404-657-2588

State Epidemiologist: Cherie L. Drenzek, DVM, MS

Office (direct line) 404-657-2609

Cherie.Drenzek@dph.ga.gov

1-866-PUB-HLTH



When it Comes to our Microbial Opponents...

